ISSN: 2642-1747

Short Communication

Copyright© John D. Scott

Human Babesiosis Caused by *Babesia odocoilei*: An Emerging Zoonosis

John D. Scott* and Catherine M. Scott

Upper Grand Tick Research Centre, 365 St. David Street South, Fergus, Ontario N1M 2L7, Canada

*Corresponding author: John D. Scott, Upper Grand Tick Research Centre, 365 St. David Street South, Fergus, Ontario N1M 2L7, Canada.

To Cite This Article: John D. Scott* and Catherine M. Scott. Human Babesiosis Caused by Babesia odocoilei: An Emerging Zoonosis. Am J Biomed Sci & Res. 2024 24(6): 667-668. AJBSR.MS.ID.003261, DOI: 10.34297/AJBSR.2024.24.003261

Received:

Movember 20, 2024; Published:

December 02, 2024

Abstract

Human babesiosis caused by *Babesia odocoilei* has an innate ability to survive in humans. *Babesia odocoilei*, a sequestered *Babesia* sp. has unique survival mechanisms to occlude capillaries and venules by forming fibrin-bonded entanglements. Immune evasion has been mastered, and establishment of persistent infections promote survival. Because *B. odocoilei* sequesters, this tick-borne zoonosis is recalcitrant to treat.

Short Communication

An emerging tick-borne zoonotic disease called human babesiosis caused by *Babesia odocoilei* is widely distributed in North America. This piroplasmid, a single-celled, red blood cell parasite causes a multitude of insidious symptoms in humans. This microscopic hemoparasite is commonly transmitted by a *B. odocoilei*-infected *Ixodes* tick, but can be transmitted by blood transfusion, organ transplantation, and congenital transmission.

Tick vectors include the western blacklegged tick, Ixodes pacificus and the blacklegged tick, Ixodes scapularis. Mammalian reservoirs are primarily cervids (i.e., white-tailed deer, Odocoileus virginianus; black-tailed deer, Odocoileus hemionus columbianus), but desert bighorn sheep, Ovis canadensis nelsoni, also harbour B. odocoilei [1]. The first-ever discovery of B. odocoilei in humans was made by Scott et al. [2]. In Canada, B. odocoilei has been detected in three tick species (i.e., Ixodes angustus, Ixodes pacificus, and Ixodes scapularis) continent wide [3]. Scott and colleagues found that the ratio of B. odocoilei to Babesia microti in I. scapularis adults is 60 to 1 [3]. Babesia odocoilei is, undeniably, the predominant Babesia sp. across North America. Based on testing 224 Ixodes ticks, researchers found that people are just as likely to contract human babesiosis as Lyme disease [3]. In fact, the Canada-wide prevalence of B. burgdorferi and B. odocoilei in *I. scapularis* adults was 40% and 36%, respectively [3]. In B.C., a person is pathologically more likely to be bitten by a *B. odocoilei*-positive *Ixodes* tick than a *B. burgdor-feri*-positive *Ixodes* tick [3,4].

Migratory songbirds play an integral role in the wide dispersal of I. scapularis larvae and nymphs. Ixodes scapularis harbour at least six tick-borne zoonotic pathogens that infect humans [4]. In the spring, northward-bound songbirds transport ticks into Canada from as far south as the neotropics and, alternatively, in the fall, southward-migrating songbirds transport ticks to southern latitudes. In Canada, the predominant pathogens that are transmitted to humans are B. burgdorferi and B. odocoilei [3]. Based on a tick-hostpathogen study, researchers detected five tick-borne zoonotic pathogens in the blood of songbirds [5]. As well, five pathogens have been detected in several studies [3,6]. Babesia odocoilei and B. burgdorferi were the most prominent pathogens [6]. When walking in forest and grassy habitat, it is just as likely for a juvenile I. scapularis to be infected with B. odocoilei (human babesiosis) as B. burgdorferi (Lyme disease) [3].

When a *B. odocoilei*-infected *I. scapularis* takes a blood meal, it spews sporozoites into the blood stream, and they quickly circulated throughout the body. Fibrinogen in the Am J Biomed Sci & Res Copyright© John D. Scott

blood converts to fibrin, and adheres to the tubular endothelium (cytoadhesion), and clogs in capillaries and post-capillary venules (sequestration) [7,8]. Pathologically fibrin encompasses both infected Red Blood Cells (iRBCs) and uninfected Red Blood Cells (uRBCs) [7,8]. These 3 components form fibrin-bonded entanglements that block capillaries and venules, especially in the brain that have the smallest capillaries. Sequestering Babesia species (i.e., B. bovis, B. canis, B. odocoilei) can complete their life cycle within fibrin-bonded entanglements and, thus, trophozoites and merozoites remains isolated from the circulatory immune system and spleen [9]. Immune evasion is ostensible. For babesial survival, iRBCs, iRBCs and fibrin form occlusions, and self-perpetuate life within these entanglements. As cytoadhesion progresses and sequestration builds (using mature babesial stages) [7], B. odocoilei has an exceptional means to avoid splenic clearance [9]. In contrast, non-sequestering Babesia (i.e., B. microti) allow the immune system, which consist of microphages, to function normally, and the spleen typically traps and destroys babesial merozoites. Subsequently, patients with B. microti are normally much easier to treat successfully.

Common Symptoms

Because fibrin-bonded entanglements occlude capillaries and venules, oxygen and nutrient are greatly reduced. As B. odocoilei propagates, a multitude of symptoms develop, including unyielding fatigue, cognitive impairment, perpetual inflammation (especially in legs at night), restless legs, brain fog, anxiety, delirium/disorientation, nightmares, profound wild dreams, disorientation, difficulty remembering, progressive dementia, dizziness, memory loss, muscle and joint stiffness, muscle and joint aches, clumsiness, poor balance, unsteady gait, bladder dysfunction, intestinal problems, constipation, sleep disturbance, insomnia, sweats (especially at night), irritability/rage/aggression, irritability, ischemia (slow blood flow), chills, heat and cold intolerance, pathogen-induced depression, air hunger, longstanding headaches, encephalopathy, and Jarisch-Herxheimer reaction (herxing). Of particular note, human babesiosis caused by B. odocoilei inflict major depression in the population. A comorbidity, such as human babesiosis caused by B. odocoilei and Lyme disease caused by B. burgdorferi s.l. can cause pain and suffering.

As *B. odocoilei* advances, patients show severe clinical signs and symptoms. Patients can become bedridden and disabled as manifestations progress. As these symptoms develop, patients can be suicidal/homicidal and, in some cases,

can have fatal outcomes [3]. As fibrin-bonded entanglements of *B. odocoilei* occlude capillaries and venules, children often become non-verbal, develop muscle weakness, and have unsteady gait. Moreover, children often have bladder and bowel dysfunctional. In particular, *B. odocoilei* is a sequestering *Babesia* that is recalcitrant to treat [3]. In contrast, *B. microti*, is a non-sequestering *Babesia* sp. that is typically less difficult to treat. However, *B. odocoilei* has developed highly effective mechanism for parasite survival in humans. Deer culls are needed to reduce human babesiosis caused by *B. odocoilei*. Tick repellants are efficacious. Since people spend considerable time in the outdoors, healthcare professionals must acquire continuing medical education in tick-borne zoonotic diseases.

Acknowledgement

None

Conflict of Interest

None.

References

- Thomford JW, Conrad PA, Boyce WM, Holman PJ, Jessop DA et al., (1993) Isolation and in vitro cultivation of *Babesia* parasites from free-ranging desert bighorn sheep (*Ovis canadensis nelsoni*) and mule deer (*Odocoil-eus hemionus*) in California. J Parasitol 79(1): 77-84.
- Scott JD, Sajid MS, Pascoe EL, Foley JE (2021) Detection of *Babesia odo-coilei* in humans with babesiosis symptoms. Diagnostics 11(6): 947.
- Scott JD, Scott CM (2024) Molecular detection of Borrelia burgdorferi sensu lato, Borrelia miyamotoi, Babesia odocoilei, Babesia microti and Anaplasma phagocytophilum in Ixodes ticks collected across Canada. J Biomed Res Environ Sci 5(1): 1321-1337.
- Scott JD, Scott CM (2023) Primary detection of the establishment of blacklegged ticks, *Ixodes scapularis*, in British Columbia, Canada. J Biomed Res Environ Sci 4(1): 935-941.
- Scott JD, McGoey E, Morales A, Pesapane RR (2022) Molecular detection of *Anaplasma phagocytophilum*, *Babesia odocoilei*, *Babesia* species and *Borrelia burgdorferi* sensu lato in songbirds. J Biomed Res Environ Sci 3(1): 1451-1459.
- Scott JD, Pesapane RR. (2021) Detection of Anaplasma phagocytophilum, Babesia odocoilei, Babesia sp., Borrelia burgdorferi sensu lato, and Hepatozoon canis in Ixodes scapularis ticks collected in eastern Canada. Pathogens 10(10): 1265.
- Wright IG (1972) An electron microscopic study of intravascular agglutination in the cerebral cortex due to *Babesia argentina* infection. Int J Parasitol 2(2): 209-205.
- Schetters TP, Kleuskens J, Scholtes N, Gorenflot A (1998) Parasite localization and dissemination in the *Babesia*-infected host. Ann Trop Med Parasitol 92(4): 513-519.
- Allred DR, Al-Khedery B (2004) Antigenic variation and cytoadhering in Babesia bovis and Plasmodium falciparum: different logics achieve the same goal. Mol Biochem Parasitol 134(1): 27-35.