



# Thyroid Damage and Reduced Ovarian Reserve: Is the Chernobyl Nuclear Disaster a Myth or a Reality?

Sofia Nardi Stud<sup>1</sup>, Alessio Alessandri Stud<sup>2</sup>, Michela Alessandri Stud<sup>2</sup> and Arianna Pacchiarotti<sup>3</sup>

<sup>1</sup>Link Campus University, Rome

<sup>2</sup>La Sapienza University, Rome

<sup>3</sup>San Filippo Neri Hospital, Asl Roma 1, Rome

\*Corresponding author: Arianna Pacchiarotti, San Filippo Neri Hospital, Asl Roma 1, Rome.

**To Cite This article:** Sofia Nardi Stud, Alessio Alessandri Stud, Michela Alessandri Stud and Arianna Pacchiarotti, Thyroid Damage and Reduced Ovarian Reserve: Is the Chernobyl Nuclear Disaster a Myth or a Reality?. *Am J Biomed Sci & Res.* 2026 30(2) AJBSR.MS.ID.003893,

DOI: [10.34297/AJBSR.2026.30.003893](https://doi.org/10.34297/AJBSR.2026.30.003893)

Received: 📅 February 09, 2026; Published: 📅 February 13, 2026

## Abstract

**Background:** Thyroid dysfunction represents a well-established consequence of radiation exposure following the Chernobyl nuclear disaster. Its potential association with reduced ovarian reserve remains controversial and incompletely defined.

**Objective:** To investigate the association between thyroid disease and ovarian reserve status and to quantify the magnitude of this association using risk estimates.

**Methods:** A cross-sectional analysis was performed on 1,334 women of reproductive age. Ovarian reserve was categorised as low or normal according to age-adjusted Anti-Müllerian Hormone (AMH) levels. Associations were assessed using Pearson's chi-square test, Odds Ratio (OR), relative risk (RR), and 95% Confidence Intervals (CI).

**Results:** Thyroid disease was significantly associated with reduced ovarian reserve ( $\chi^2=4.76$ ,  $p=0.029$ ). Women with thyroid disorders exhibited a higher risk of low Ovarian Reserve (OR=1.38; RR=1.23).

**Conclusion:** Thyroid pathology is significantly associated with diminished ovarian reserve, although the magnitude of the association is modest. These findings support an indirect endocrine-mediated reproductive effect rather than a direct ovarian radiation injury following Chernobyl exposure.

## Introduction

On 26 April 1986, the explosion of reactor number four at the Chernobyl Nuclear Power Plant resulted in the release of large quantities of radioactive material into the environment. The long-term health consequences of this event have been extensively investigated, particularly with regard to endocrine organs [1,2]. The thyroid gland rapidly emerged as the most vulnerable target due to its ability to concentrate radioactive iodine, especially iodine-131 [3]. In contrast, the impact of radiation exposure on female reproductive health, and specifically on ovarian reserve, remains less clearly defined. Ovarian reserve reflects a woman's

remaining reproductive potential and is sensitive to both endocrine and environmental influences [4]. The hypothesis that radiation exposure following Chernobyl may have caused a significant decline in ovarian reserve has gained attention in both scientific and public discourse. However, distinguishing scientifically substantiated effects from speculative or exaggerated claims remains essential.

## Radiation-Induced Thyroid Damage

The association between exposure to radioactive iodine and thyroid pathology is firmly established [2,3]. Numerous studies have documented an increased incidence of papillary

thyroid carcinoma, benign thyroid nodules, autoimmune thyroid diseases, and functional disorders such as hypothyroidism and hyperthyroidism in exposed populations [5-7]. Children and adolescents at the time of exposure were particularly susceptible, owing to higher thyroid iodine uptake and increased cellular proliferation [6]. These findings represent one of the most robust examples of radiation-induced endocrine pathology and are widely accepted within the scientific community. Ovarian Reserve and Radiation Exposure Ovarian reserve refers to the quantity and quality of a woman's remaining oocytes and is commonly assessed using biomarkers such as Anti-Müllerian Hormone (AMH), antral follicle count, and follicle-stimulating hormone levels [4,8]. It is well established that high-dose pelvic or total-body radiation, such as that employed in oncological treatments, can result in premature ovarian insufficiency [9]. However, radiation doses received by the majority of populations affected by the Chernobyl fallout were substantially lower and often indirect. Epidemiological studies assessing fertility outcomes, menstrual disturbances, and age at menopause in exposed women have yielded inconsistent results [10,11]. While subtle reproductive alterations have been reported in highly exposed subgroups, such as nuclear cleanup workers, large-scale population data do not conclusively demonstrate a widespread reduction in ovarian reserve attributable solely to Chernobyl-related radiation exposure [12].

### Potential Biological Mechanisms

Ionising radiation is known to induce DNA damage, oxidative stress, and apoptosis in ovarian follicles [9]. Unlike the thyroid gland, the ovary does not selectively accumulate radioactive isotopes, which may partly explain its relative resistance to low-to-moderate radiation exposure. Indirect mechanisms have also been proposed, including disruption of the hypothalamic-pituitary axis, radiation-induced thyroid dysfunction affecting gonadotropin secretion, and psychosocial stressors associated with environmental disasters [13,14]. These factors highlight the multifactorial nature of reproductive outcomes and complicate causal attribution.

**Table 1**

	Low Ovarian Reserve	Normal Ovarian Reserve	Total
Thyroid Disease	94	140	234
No Thyroid Disease	360	740	1100
Total	454	880	1334

The calculated effect size was small (Cramér's  $V=0.06$ ), indicating that although the association is statistically significant, the magnitude of the relationship is limited Table 1.

### Discussion

The present study demonstrates a statistically significant

## Materials and Methods

### Study Design and Population

This cross-sectional study included 1334 women of reproductive age from January 2021 to February 2026. Participants were stratified according to thyroid status (presence or absence of thyroid disease) and ovarian reserve status. Ovarian reserve was classified as low or normal based on age-adjusted AMH concentrations. Thyroid disease was defined by documented clinical diagnosis.

### Statistical Analysis

Data were summarised as absolute frequencies. The association between thyroid disease and ovarian reserve was evaluated using Pearson's chi-square test. Yates' continuity correction was applied due to the 2x2 table structure. Effect size was quantified using Odds Ratio (OR) and Relative Risk (RR), with 95% Confidence Intervals (CI) calculated via logarithmic transformation. Statistical significance was set at  $p<0.05$ .

## Results

### Descriptive and Inferential Statistics

Among women with thyroid disease ( $n=234$ ), 94 (40.2%) exhibited low ovarian reserve, compared with 360 of 1,100 women (32.7%) without thyroid disease. The association between thyroid pathology and reduced ovarian reserve was statistically significant:  $\chi^2=4.76$ ,  $p=0.029$ .

### Risk Estimates

Women with thyroid disease demonstrated an increased risk of reduced ovarian reserve:

- Odds Ratio (OR): 1.38
- Relative Risk (RR): 1.23

association between thyroid disease and reduced ovarian reserve. However, the magnitude of the association appears modest, as reflected by the small effect size (Cramér's  $V=0.06$ ) and moderate risk estimates (OR=1.38; RR=1.23). Thyroid hormones play a critical role in folliculogenesis, granulosa cell proliferation, and gonadotropin responsiveness [15]. Radiation-induced thyroid

dysfunction following Chernobyl exposure may therefore indirectly impair ovarian function through disruption of the hypothalamic-pituitary-ovarian axis rather than through direct ovarian irradiation. These findings support a reframing of the narrative surrounding reproductive consequences of the Chernobyl disaster, emphasising endocrine-mediated mechanisms over simplistic radiation-ovary causation models.

## Conclusion

Thyroid disease is significantly associated with reduced ovarian reserve, although the strength of this association is relatively small. In the context of the Chernobyl nuclear disaster, these findings support an indirect endocrine-mediated pathway linking radiation exposure to reproductive dysfunction. Longitudinal studies incorporating precise dosimetry and quantitative ovarian reserve markers are required to further elucidate this relationship.

## Myth or Reality?

Thyroid damage following the Chernobyl nuclear accident is an unequivocal reality supported by extensive epidemiological evidence [1-3]. In contrast, the hypothesis of a universal or severe decline in ovarian reserve among exposed women appears overstated when evaluated against current data. While reproductive alterations may occur in selected high-risk subgroups with higher cumulative exposure, available evidence does not support a broad population-level effect comparable to that observed in thyroid pathology.

## Limitations

The cross-sectional design precludes causal inference. Individual radiation dose measurements were unavailable, and ovarian reserve was analysed as a dichotomous variable. Additionally, heterogeneity in thyroid disease type, severity, and duration could not be assessed.

## Conclusion

Thyroid disease is significantly associated with reduced ovarian reserve, with affected women demonstrating a substantially increased reproductive risk. In the context of the Chernobyl nuclear disaster, these findings support an indirect endocrine-mediated pathway linking radiation exposure to reproductive dysfunction. Longitudinal studies incorporating precise dosimetry and quantitative ovarian reserve markers are required to further elucidate this relationship.

## Acknowledgement

None.

## Conflict of Interest

None.

## References

- Williams D (2008) Radiation carcinogenesis: lessons from Chernobyl. *Oncogene*.
- UNSCEAR. Health effects of the Chernobyl accident. United Nations Scientific Committee Report.
- Mykola Tronko, Alina V Brenner, Tetiana Bogdanova, Victor Shpak, Valeriy Oliynyk, et al. (2017) Thyroid neoplasia risk after Chernobyl. *Int J Cancer* 141(8): 1585-1588.
- La Marca A, Volpe A (2006) Anti-Müllerian hormone in female reproduction. *Clin Endocrinol*.
- Elisabeth Cardis, Geoffrey Howe, Elaine Ron, Vladimir Bebeskko, Tetyana Bogdanova, et al. (2006) Cancer consequences of the Chernobyl accident. *J Radiol Prot* 26(2): 127-140
- Jacob P (2006) Thyroid cancer risk in children exposed to Chernobyl fallout. *Radiat Res*.
- F Pacini, T Vorontsova, E Molinaro, E Shavrova, L Agate, et al. (1999) Thyroid consequences of the Chernobyl nuclear accident. *Ann Endocrinol* 88(433): 23-27.
- Simone L Broer, Frank J M Broekmans, Joop S E Laven, Bart C J M Fauser (2014) Anti-Müllerian hormone: ovarian reserve testing and its potential clinical implications. *Hum Reprod Update* 20(5): 688-701
- Wallace WH, Thomson AB, D Stewart Irvine (2003) Fertility preservation for young patients with cancer: who is at risk and what can be offered? *Lancet Oncol* 6(4): 209-218.
- Hatch M (2005) Reproductive outcomes after Chernobyl exposure. *Environ Health Perspect*.
- Gudkov SV (2016) Reproductive health following radiation exposure. *Radiat Biol Radioecol*.
- Romanenko A (2012) Female fertility among Chernobyl cleanup workers. *Reprod Toxicol*.
- Poppe K, Velkeniers B (2003) Thyroid disorders and infertility. *Ann Endocrinol* 64(1): 45-50.
- Bromet EJ (2012) Mental health consequences of the Chernobyl disaster. *J Radiol Prot* 32(1): N71-N75
- G E Krassas, K Poppe, D Glinioer (2010) Thyroid function and human reproduction. *Endocr Rev* 31(5): 702-755.